



CROWN PREPARATION AFTERCARE

You may experience some mild tenderness and swelling at the site of the injection. The resulting numbness/tingling is typically short acting but may be prolonged in some individuals.

Please take care while you are still numb to not accidentally bite your e.g. cheek/lip/tongue. Please avoid very hot/cold foods/drinks until you regain sensation around the area.

Sensitivity is a common reaction following any complex dental procedure. This is generally short-lived and will resolve with time.

It is very important you notify your dentist of any symptoms by the day of fitting the final crown

TEMPORARY CROWN

You have been given a temporary crown, while the crown is being made. This is an important part of your treatment as it will not only protect your tooth, but also stops the neighbouring and opposing teeth from moving.

The temporary crown is not an indication of the colour and polish of the final product.

The temporary is stuck in loosely by design. Care must be taken to ensure it does not dislodge prematurely. Please avoid hard/crunchy/sticky textures and use the other side if possible.

Flossing around the temporary crown is crucial to maintain the health of the gum around the tooth, but please do so with care as instructed by your dentist

If the temporary crown falls off or is lost, please call us immediately to make an appointment to have it fitted back on, or a new one made. Bringing the temporary crown with you to place back on is not crucial, but it can shorten your appointment time.

Please note that if the tooth is left too long (generally more than 4 days) without the temporary crown, the final crown may not fit and may have to be redone.